

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 635 507 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
15.09.1999 Bulletin 1999/37

(21) Application number: 94305194.6

(22) Date of filing: 15.07.1994

(51) Int Cl.⁶: C07D 487/04, A61K 31/495,
C07D 498/04, C07D 513/04
// (C07D487/04, 239:00,
235:00),
(C07D487/04, 249:00, 239:00),
(C07D487/04, 241:00, 239:00)

(54) **Tricyclic derivatives and their use as anti-cancer agents**

Tricyclische Derivate und ihre Anwendung als Krebsmittel

Dérivés tricyclique et leur utilisation comme agents anti-cancer

(84) Designated Contracting States:
AT BE CH DE DK ES FR GB GR IE IT LI LU MC NL
PT SE
Designated Extension States:
LT SI

(30) Priority: 19.07.1993 GB 9314884

(43) Date of publication of application:
25.01.1995 Bulletin 1995/04

(73) Proprietor: ZENECA LIMITED
London W1Y 6LN (GB)

(72) Inventor: Barker, Andrew John
Cheshire SK10 4TG (GB)

(74) Representative: Gilles, David Eric et al
AstraZeneca PLC
Global Intellectual Property
Meriside
Alderley Park
Macclesfield, Cheshire SK10 4TG (GB)

(56) References cited:
EP-A- 0 520 722 EP-A- 0 542 797

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 0 635 507 B1

Description

[0001] The invention relates to tricyclic derivatives, or pharmaceutically-acceptable salts thereof, which possess anti-cancer activity and are accordingly useful in methods of treatment of cancer in the human or animal body. The invention also relates to processes for the manufacture of said tricyclic derivatives, to pharmaceutical compositions containing them and to their use in the manufacture of medicaments of use in the production of an anti-cancer effect in a warm-blooded animal such as man.

[0002] Many of the current treatment regimes for cancer utilise compounds which inhibit DNA synthesis. Such compounds are toxic to cells generally but their toxic effect on the rapidly dividing tumour cells can be beneficial. Alternative approaches to anti-cancer agents which act by mechanisms other than the inhibition of DNA synthesis have the potential to display enhanced selectivity of action against cancer cells.

[0003] In recent years it has been discovered that a cell may become cancerous by virtue of the transformation of a portion of its DNA into an oncogene i.e. a gene which, on activation, leads to the formation of malignant tumour cells (Bradshaw, *Mutagenesis*, 1986, **1**, 91). Several such oncogenes give rise to the production of peptides which are receptors for growth factors. The growth factor receptor complex subsequently leads to an increase in cell proliferation. It is known, for example, that several oncogenes encode tyrosine kinase enzymes and that certain growth factor receptors are also tyrosine kinase enzymes (Yarden *et al.*, *Ann. Rev. Biochem.*, 1988, **57**, 443; Larsen *et al.* *Ann. Reports in Med. Chem.* 1989, Chpt. 13).

[0004] Receptor tyrosine kinases are important in the transmission of biochemical signals which initiate cell replication. They are large enzymes which span the cell membrane and possess an extracellular binding domain for growth factors such as epidermal growth factor and an intracellular portion which functions as a kinase to phosphorylate tyrosine amino acids in proteins and hence to influence cell proliferation. It is known that such kinases are frequently present in common human cancers such as breast cancer (Sainsbury *et al.*, *Brit. J. Cancer*, 1988, **58**, 458; Guerin *et al.*, *Oncogene Res.*, 1988, **3**, 21), gastrointestinal cancer such as colon, rectal or stomach cancer (Bolen *et al.*, *Oncogene Res.*, 1987, **1**, 149), leukaemia (Konaka *et al.*, *Cell*, 1984, **37**, 1035) and ovarian, bronchial or pancreatic cancer (European Patent Specification No. 0400586). As further human tumour tissues are tested for receptor tyrosine kinase activity it is expected that its widespread prevalence will be established in further cancers such as thyroid and uterine cancer. It is also known that tyrosine kinase activity is rarely detected in normal cells whereas it is more frequently detectable in malignant cells (Hunter, *Cell*, 1987, **50**, 823). It has been shown more recently (W J Gullick, *Brit. Med. Bull.*, 1991, **47**, 87) that epidermal growth factor receptor which possesses tyrosine kinase activity is overexpressed in many human cancers such as brain, lung squamous cell, bladder, gastric, breast, head and neck, oesophageal, gynaecological and thyroid tumours.

[0005] Accordingly it has been recognised that an inhibitor of receptor tyrosine kinase should be of value as a selective inhibitor of the growth of mammalian cancer cells (Yaish *et al.*, *Science*, 1988, **242**, 933). Support for this view is provided by the demonstration that erbstatin, a receptor tyrosine kinase inhibitor, specifically attenuates the growth in athymic nude mice of a transplanted human mammary carcinoma which expresses epidermal growth factor (EGF) receptor tyrosine kinase but is without effect on the growth of another carcinoma which does not express EGF receptor tyrosine kinase (Toi *et al.*, *Eur. J. Cancer Clin. Oncol.*, 1990, **26**, 722.) Various derivatives of styrene are also stated to possess tyrosine kinase inhibitory properties (European Patent Application Nos. 0211363, 0304493 and 0322738) and to be of use as anti-tumour agents. The *in vivo* inhibitory effect of two such styrene derivatives has been demonstrated against the growth of human squamous cell carcinoma inoculated into nude mice (Yoneda *et al.*, *Cancer Research*, 1991, **51**, 4430). Accordingly it has been indicated that receptor tyrosine kinase inhibitors will prove to be useful in the treatment of a variety of human cancers. Various known tyrosine kinase inhibitors are disclosed in a more recent review by T R Burke Jr. (*Drugs of the Future*, 1992, **17**, 119).

[0006] We have now found that certain tricyclic derivatives which incorporate a quinazoline ring possess anti-cancer properties which are believed to arise from their receptor tyrosine kinase inhibitory properties.

[0007] It is known from the patent application WO 92/20642 that certain aryl and heteroaryl compounds inhibit receptor tyrosine kinase. There is the disclosure of certain quinazoline derivatives but no mention is made of 4-anilinoquinazoline derivatives.

[0008] It is also known from European Patent Application No. 92305703.8 (publication no. 0 520 722) that certain 4-anilinoquinazoline derivatives which are unsubstituted at positions 5 to 8 of the quinazoline ring or which bear a halogeno, trifluoromethyl or nitro substituent at one of those positions are useful as inhibitors of receptor tyrosine kinase.

[0009] According to the present invention there is provided a tricyclic derivative of the formula I (set out hereinafter) wherein R¹ and R² together form a group of the formula -N=CH-NH-, -N=CH-O-, -N=CH-S-, -N=N-NH-, -NH-N=CH-, -NH-CH=CH-, -NH-CO-NH-, -NH-CO-O-, -NH-CO-S-, -NH-NH-CO-, -N=CH-CH=CH-, -N=N-CH=CH-, -N=CH-CH=N-, -NH-CO-CH=CH- or -N=CH-CO-NH- (with in each case a nitrogen atom being located at the 6-position of the quinazoline ring) and the 5- or 6-membered ring so formed may optionally bear one or two substituents, any substituent on an available nitrogen atom being selected from (1-4C)alkyl, (3-4C)alkenyl, (3-4C)alkynyl, halogeno-

(1-4C)alkyl, hydroxy-(1-4C)alkyl, (2-4C)alkanoyloxy-(1-4C)alkyl, (1-4C)alkoxy-(1-4C)alkyl, cyano-(1-4C)alkyl, amino-(1-4C)alkyl, (1-4C)alkylamino-(1-4C)alkyl and di-[(1-4C)alkyl]amino-(1-4C)alkyl, and any substituent on an available carbon atom being selected from halogeno, amino, hydroxy, carbamoyl, cyano, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, (1-4C)alkoxy, (1-4C)alkylthio, (1-4C)alkylsulphinyl, (1-4C)alkylsulphonyl, (1-4C)alkylamino, di-[(1-4C)alkyl]amino, (2-4C)alkanoyl, N-(1-4C)alkylcarbamoyl, N,N-di-[(1-4C)alkyl]carbamoyl, halogeno-(1-4C)alkyl, hydroxy-(1-4C)alkyl, (2-4C)alkanoyloxy-(1-4C)alkyl, (1-4C)alkoxy-(1-4C)alkyl, cyano-(1-4C)alkyl, amino-(1-4C)alkyl, (1-4C)alkylamino-(1-4C)alkyl and di-[(1-4C)alkyl]amino-(1-4C)alkyl; and

m is the integer 1, 2 or 3 and each R³ is independently hydrogen, halogeno, trifluoromethyl, hydroxy, amino, nitro, cyano, (1-4C)alkyl, (1-4C)alkoxy, (1-4C)alkylamino, di-[(1-4C)alkyl]amino or (2-4C)alkanoylamino;

or a pharmaceutically-acceptable salt thereof.

[0010] The chemical formulae referred to herein by Roman numerals are set out for convenience on a separate sheet hereinafter. In this specification the term "alkyl" includes both straight and branched chain alkyl groups but references to individual alkyl groups such as "propyl" are specific for the straight chain version only. An analogous convention applies to other generic terms.

[0011] Within the present invention it is to be understood that a quinazoline of the formula I may exhibit the phenomenon of tautomerism and that the formulae drawings within this specification can represent only one of the possible tautomeric forms. It is to be understood that the invention encompasses any tautomeric form which possesses anti-cancer activity and is not to be limited merely to any one tautomeric form utilised within the formulae drawings.

[0012] The quinazolines of the formula I are unsubstituted at the 2-, 5- and 8-positions.

[0013] It is also to be understood that certain quinazolines of the formula I can exist in solvated as well as unsolvated forms such as, for example, hydrated forms. It is to be understood that the invention encompasses all such solvated forms which possess anti-cancer activity.

[0014] According to a further aspect of the present invention there is provided a tricyclic derivative of the formula I wherein R¹ and R² together form a group of the formula -N=CH-NH-, -N=CH-O-, -N=CH-S-, -N=N-NH-, -NH-N=CH-, -NH-CH=CH-, -NH-CO-NH-, -NH-CO-O-, -NH-CO-S-, -NH-NH-CO-, -N=CH-CH=CH-, -N=N-CH=CH-, -N=CH-N=CH-, -N=CH-CH=N- or -NH-CO-CH=CH- (with in each case a nitrogen atom being located at the 6-position of the quinazoline ring) and the 5- or 6-membered ring so formed may optionally bear one or two substituents, any substituent on an available nitrogen atom being selected from (1-4C)alkyl, (3-4C)alkenyl, (3-4C)alkynyl, halogeno-(1-4C)alkyl, hydroxy-(1-4C)alkyl, (2-4C)alkanoyloxy-(1-4C)alkyl, (1-4C)alkoxy-(1-4C)alkyl, cyano-(1-4C)alkyl, amino-(1-4C)alkyl, (1-4C)alkylamino-(1-4C)alkyl and di-[(1-4C)alkyl]amino-(1-4C)alkyl, and any substituent on an available carbon atom being selected from halogeno, amino, carbamoyl, cyano, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, (1-4C)alkoxy, (1-4C)alkylthio, (1-4C)alkylsulphinyl, (1-4C)alkylsulphonyl, (1-4C)alkylamino, di-[(1-4C)alkyl]amino, (2-4C)alkanoyl, N-(1-4C)alkylcarbamoyl, N,N-di-[(1-4C)alkyl]carbamoyl, halogeno-(1-4C)alkyl, hydroxy-(1-4C)alkyl, (2-4C)alkanoyloxy-(1-4C)alkyl, (1-4C)alkoxy-(1-4C)alkyl, cyano-(1-4C)alkyl, amino-(1-4C)alkyl, (1-4C)alkylamino-(1-4C)alkyl and di-[(1-4C)alkyl]amino-(1-4C)alkyl; and

m is the integer 1, 2 or 3 and each R³ is independently hydrogen, halogeno, trifluoromethyl, hydroxy, amino, nitro, cyano, (1-4C)alkyl, (1-4C)alkoxy, (1-4C)alkylamino, di-[(1-4C)alkyl]amino or (2-4C)alkanoylamino;

or a pharmaceutically-acceptable salt thereof.

[0015] Suitable values for the generic radicals referred to above include those set out below.

[0016] Suitable values for each substituent which may be present on the ring involving R¹ and R², or for each R³ substituent which may be present include, for example:-

for halogeno	fluoro, chloro, bromo and iodo;
for (1-4C)alkyl	methyl, ethyl, propyl, isopropyl, butyl, isobutyl, <u>sec</u> -butyl and <u>tert</u> -butyl;
for (3-4C)alkenyl	allyl and but-2-enyl;
for (2-4C)alkenyl	vinyl, allyl and but-2-enyl;
for (3-4C)alkynyl	prop-2-ynyl and but-2-ynyl;
for (2-4C)alkynyl	ethynyl, prop-2-ynyl and but-2-ynyl;
for (1-4C)alkoxy	methoxy, ethoxy, propoxy, isopropoxy and butoxy;
for (1-4C)alkylthio	methylthio, ethylthio and propylthio;
for (1-4C)alkylsulphinyl	methylsulphinyl, ethylsulphinyl and propylsulphinyl;
for (1-4C)alkylsulphonyl	methylsulphonyl, ethylsulphonyl and propylsulphonyl;
for (1-4C)alkylamino	methylamino, ethylamino and propylamino;
for di-[(1-4C)alkyl]amino	dimethylamino, <u>N</u> -ethyl- <u>N</u> -methylamino, diethylamino, <u>N</u> -methyl- <u>N</u> -propylamino and dipropylamino;
for (2-4C)alkanoylamino	acetamido, propionamido and butyramido;

(continued)

for (2-4C)alkanoyl	acetyl, propionyl and butyryl;
for N-(1-4C)alkylcarbamoyl	N-methylcarbamoyl, N-ethylcarbamoyl and N-propylcarbamoyl;
for N,N-di-[(1-4C)alkyl]carbamoyl	N,N-dimethylcarbamoyl, N-ethyl-N-methylcarbamoyl and N,N-diethylcarbamoyl;
for halogeno-(1-4C)alkyl	fluoromethyl, chloromethyl, bromomethyl, difluoromethyl, dichloromethyl, dibromomethyl, 2-fluoroethyl, 2,2-difluoroethyl, 2,2,2-trifluoroethyl, 2-chloroethyl, 2-bromoethyl and trifluoromethyl;
for hydroxy-(1-4C)alkyl	hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl and 3-hydroxypropyl;
for (2-4C)alkanoyloxy-(1-4C)alkyl	acetoxymethyl, propionyloxymethyl, butyryloxymethyl, 2-acetoxyethyl and 3-acetoxypentyl;
for (1-4C)alkoxy-(1-4C)alkyl	methoxymethyl, ethoxymethyl, 1-methoxyethyl, 2-methoxyethyl, 2-ethoxyethyl and 3-methoxypropyl;
for cyano-(1-4C)alkyl	cyanomethyl, 1-cyanoethyl, 2-cyanoethyl and 3-cyanopropyl;
for amino-(1-4C)alkyl	aminomethyl, 1-aminoethyl, 2-aminoethyl and 3-aminopropyl;
for (1-4C)alkylamino-(1-4C)alkyl	methylaminomethyl, ethylaminomethyl, 1-methylaminoethyl, 2-methylaminoethyl, 2-ethylaminoethyl and 3-methylaminopropyl;
for di-[(1-4C)alkyl]amino(1-4C)alkyl	dimethylaminomethyl, diethylaminomethyl, 1-dimethylaminoethyl, 2-dimethylaminoethyl and 3-dimethylaminopropyl.

[0017] A suitable pharmaceutically-acceptable salt of a tricyclic derivative of the invention is, for example, an acid-addition salt of a tricyclic derivative of the invention which is sufficiently basic, for example, an acid-addition salt with, for example, an inorganic or organic acid, for example hydrochloric, hydrobromic, sulphuric, phosphoric, trifluoroacetic, citric or maleic acid. In addition a suitable pharmaceutically-acceptable salt of a tricyclic derivative of the invention which is sufficiently acidic is an alkali metal salt, for example a sodium or potassium salt, an alkaline earth metal salt, for example a calcium or magnesium salt, an ammonium salt or a salt with an organic base which affords a physiologically-acceptable cation, for example a salt with methylamine, dimethylamine, trimethylamine, piperidine, morpholine or tris-(2-hydroxyethyl)amine.

[0018] Particular novel compounds of the invention include, for example, tricyclic derivatives of the formula I, or pharmaceutically-acceptable salts thereof wherein:-

(a) the optionally-substituted tricyclic ring defined by the linking of the R¹ and R² substituents on the quinazoline of the formula I is selected from 3H-imidazo[4,5-g]quinazolin-8-yl, oxazolo[4,5-g]quinazolin-8-yl, thiazolo[4,5-g]quinazolin-8-yl, 3H-[1,2,3]triazolo[4,5-g]quinazolin-8-yl, 1H-pyrazolo[3,4-g]quinazolin-8-yl, 6H-pyrrolo[2,3-g]quinazolin-4-yl, 2-oxo-1,2-dihydro-3H-imidazo[4,5-g]quinazolin-8-yl, 2-oxo-1,2-dihydrooxazolo[4,5-g]quinazolin-8-yl, 2-oxo-1,2-dihydrothiazolo[4,5-g]quinazolin-8-yl and 3-oxo-2,3-dihydro-1H-pyrazolo[3,4-g]quinazolin-8-yl; and m and R³ have any of the meanings defined hereinbefore;

(b) the optionally-substituted tricyclic ring defined by the linking of the R¹ and R² substituents on the quinazoline of the formula I is selected from pyrido[2,3-g]quinazolin-4-yl, pyrimidino[4,5-g]cinnolin-9-yl, pyrimidino[4,5-g]quinazolin-4-yl, pyrazino[2,3-g]quinazolin-4-yl and 7-oxo-6,7-dihydropyrido[2,3-g]quinazolin-4-yl; and m and R³ have any of the meanings defined hereinbefore; or

(c) the optionally-substituted tricyclic ring defined by the linking of the R¹ and R² substituents on the quinazoline of the formula I is selected from pyrazino[2,3-g]quinazolin-4-yl and 8-oxo-8,9-dihydropyrazino[2,3-g]quinazolin-4-yl; and m and R³ have any of the meanings defined hereinbefore.

[0019] A further particular compound of the invention is a tricyclic derivative of the formula I

wherein the 6,6,5-tricyclic ring defined by the linking of the groups R¹ and R² is selected from 3H-imidazo[4,5-g]quinazolin-8-yl, oxazolo[4,5-g]quinazolin-8-yl, thiazolo[4,5-g]quinazolin-8-yl, 3H-[1,2,3]triazolo[4,5-g]quinazolin-8-yl, 1H-pyrazolo[3,4-g]quinazolin-8-yl, 6H-pyrrolo[2,3-g]quinazolin-4-yl, 2-oxo-1,2-dihydro-3H-imidazo[4,5-g]quinazolin-8-yl, 2-oxo-1,2-dihydrooxazolo[4,5-g]quinazolin-8-yl, 2-oxo-1,2-dihydrothiazolo[4,5-g]quinazolin-8-yl and 3-oxo-2,3-dihydro-1H-pyrazolo[3,4-g]quinazolin-8-yl, and the 5-membered ring involving R¹ and R² may optionally bear one or two substituents, any substituent on an available nitrogen atom being selected from methyl, ethyl, propyl, allyl, prop-2-ynyl, 2,2,2-trifluoroethyl, 2-hydroxyethyl, 3-hydroxypropyl, 2-acetoxyethyl, 2-methoxyethyl, 3-methoxypropyl, cyanomethyl, 2-cyanoethyl, 2-aminoethyl, 2-methylaminoethyl and 2-dimethylaminoethyl, and any substituent on an available carbon atom being selected from fluoro, chloro, amino, carbamoyl, cyano, methyl, ethyl, propyl, vinyl, allyl, ethynyl, prop-2-ynyl, methoxy, ethoxy, propoxy, methylthio, methylsulphinyl, methylsulphonyl, methylamino, dimethylamino, acetyl, N-meth-

ylcarbamoyl, *N,N*-dimethylcarbamoyl, fluoromethyl, difluoromethyl, trifluoromethyl, 2,2,2-trifluoroethyl, hydroxymethyl, 2-hydroxyethyl, acetoxymethyl, 2-acetoxyethyl, methoxymethyl, 2-methoxyethyl, cyanomethyl, 2-cyanoethyl, aminomethyl, 2-aminoethyl, methylaminomethyl, 2-methylaminoethyl, dimethylaminomethyl and 2-dimethylaminoethyl; and *m* is the integer 1, 2 or 3 and each *R*³ is independently hydrogen, fluoro, chloro, bromo, trifluoromethyl, hydroxy, amino, nitro, cyano, methyl, ethyl, methoxy, methylamino, dimethylamino or acetamido; or a pharmaceutically-acceptable salt thereof.

[0020] A further particular compound of the invention is a tricyclic derivative of the formula I wherein the 6,6,6-tricyclic ring defined by the linking of the groups *R*¹ and *R*² is selected from pyrido[2,3-*g*]quinazolin-4-yl, pyrimidino[4,5-*g*]cinnolin-9-yl, pyrimidino[4,5-*g*]quinazolin-4-yl, pyrazino[2,3-*g*]quinazolin-4-yl, 7-oxo-6,7-dihydro-pyrido[2,3-*g*]quinazolin-4-yl and 8-oxo-8,9-dihydropyrazino[2,3-*g*]quinazolin-4-yl,

and the 6-membered ring involving *R*¹ and *R*² may optionally bear one or two substituents, any substituent on an available nitrogen atom being selected from methyl, ethyl and propyl, and any substituent on an available carbon atom being selected from fluoro, chloro, hydroxy, carbamoyl, cyano, methyl, methoxy, ethoxy, *N*-methylcarbamoyl, *N,N*-dimethylcarbamoyl, trifluoromethyl and 2,2,2-trifluoroethyl; and *m* is the integer 1 or 2 and each *R*³ is independently hydrogen, fluoro, chloro, bromo, trifluoromethyl, nitro, cyano, methyl or ethyl; or a pharmaceutically-acceptable salt thereof.

[0021] A preferred compound of the invention is a tricyclic derivative of the formula I wherein the 6,6,5-tricyclic ring defined by the linking of the groups *R*¹ and *R*² is selected from 3-methyl-3*H*-imidazo[4,5-*g*]quinazolin-8-yl, 3-methyl-3*H*-[1,2,3]triazolo[4,5-*g*]quinazolin-8-yl and 3-methyl-2-oxo-1,2-dihydro-3*H*-imidazo[4,5-*g*]quinazolin-8-yl; and

(*R*³)_{*m*} is 3'-methyl, 3'-chloro or 3'-chloro-4'-fluoro; or a pharmaceutically-acceptable salt thereof.

[0022] A further preferred compound of the invention is a tricyclic derivative of the formula I wherein the 6,6,5-tricyclic ring defined by the linking of the groups *R*¹ and *R*² is selected from 3-methyl-3*H*-imidazo[4,5-*g*]quinazolin-8-yl, 3-methyl-2-oxo-1,2-dihydro-3*H*-imidazo[4,5-*g*]quinazolin-8-yl and 3-methyl-2-trifluoromethyl-3*H*-imidazo[4,5-*g*]quinazolin-8-yl; and

(*R*³)_{*m*} is 3'-methyl, 3'-chloro or 3'-chloro-4'-fluoro; or a pharmaceutically-acceptable salt thereof.

[0023] A further preferred compound of the invention is a tricyclic derivative of the formula I wherein the 6,6,6-tricyclic ring defined by the linking of the groups *R*¹ and *R*² is selected from 7,8-dimethylpyrazino[2,3-*g*]quinazolin-4-yl and 7-hydroxy-9-methyl-8-oxo-8,9-dihydropyrazino[2,3-*g*]quinazolin-4-yl; and

(*R*³)_{*m*} is 3'-methyl, 3'-chloro or 3'-chloro-4'-fluoro; or a pharmaceutically-acceptable salt thereof.

[0024] A specific preferred compound of the invention is the following tricyclic derivative of the formula I:-

3-methyl-8-(3'-methylanilino)-3*H*-imidazo[4,5-*g*]quinazoline, 3-methyl-8-(3'-methylanilino)-1,2-dihydro-3*H*-imidazo[4,5-*g*]quinazolin-2-one or 8-(3'-chloro-4'-fluoroanilino)-3-methyl-2-trifluoromethyl-3*H*-imidazo[4,5-*g*]quinazoline; or a pharmaceutically-acceptable salt thereof.

[0025] A further specific preferred compound of the invention is the following tricyclic derivative of the formula I:-

4-(3'-chloro-4'-fluoroanilino)-7,8-dimethylpyrazino[2,3-*g*]quinazoline or 4-(3'-chloro-4'-fluoroanilino)-7-hydroxy-9-methyl-8-oxo-8,9-dihydro-pyrazino[2,3-*g*]quinazoline; or a pharmaceutically-acceptable salt thereof.

[0026] A tricyclic derivative of the formula I, or a pharmaceutically-acceptable salt thereof, may be prepared by any process known to be applicable to the preparation of chemically-related compounds. A suitable process is, for example, illustrated by that used in European Patent Application No. 0 520 722. Such processes, when used to prepare a tricyclic derivative of the formula I, or a pharmaceutically-acceptable salt thereof, are provided as a further feature of the invention and are illustrated by the following representative examples in which, unless otherwise stated, *R*¹, *R*², *R*³ and *m* have any of the meanings defined hereinbefore for a tricyclic derivative of the formula I. Necessary starting materials may be obtained by standard procedures of organic chemistry. The preparation of such starting materials is described within the accompanying non-limiting Examples. Alternatively necessary starting materials are obtainable by analogous procedures to those illustrated which are within the ordinary skill of an organic chemist.

(a) The reaction, conveniently in the presence of a suitable base, of a quinazoline of the formula II (set out hereinafter), wherein *Z* is a displaceable group, with an aniline of the formula III.

A suitable displaceable group *Z* is, for example, a halogeno, alkoxy, aryloxy or sulphonyloxy group, for example a chloro, bromo, methoxy, phenoxy, methanesulphonyloxy or toluene-*p*-sulphonyloxy group.

A suitable base is, for example, an organic amine base such as, for example, pyridine, 2,6-lutidine, collidine, 4-dimethylaminopyridine, triethylamine, morpholine, *N*-methylmorpholine or diazabicyclo[5.4.0]undec-7-ene, or,

for example, an alkali or alkaline earth metal carbonate or hydroxide, for example sodium carbonate, potassium carbonate, calcium carbonate, sodium hydroxide or potassium hydroxide. Alternatively a suitable base is, for example, an alkali metal or alkaline earth metal amide, for example sodium amide or sodium bis(trimethylsilyl)amide.

The reaction is preferably carried out in the presence of a suitable inert solvent or diluent, for example an alkanol or ester such as methanol, ethanol, isopropanol or ethyl acetate, a halogenated solvent such as methylene chloride, chloroform or carbon tetrachloride, an ether such as tetrahydrofuran or 1,4-dioxan, an aromatic solvent such as toluene, or a dipolar aprotic solvent such as *N,N*-dimethylformamide, *N,N*-dimethylacetamide, *N*-methylpyrrolidin-2-one or dimethylsulphoxide. The reaction is conveniently carried out at a temperature in the range, for example, 10 to 150°C, preferably in the range 20 to 80°C.

The quinazoline derivative of the formula I may be obtained from this process in the form of the free base or alternatively it may be obtained in the form of a salt with the acid of the formula H-Z wherein Z has the meaning defined hereinbefore. When it is desired to obtain the free base from the salt, the salt may be treated with a suitable base as defined hereinbefore using a conventional procedure.

(b) For the production of those optionally-substituted compounds of the formula I wherein R¹ and R² together form a group of the formula -N=CH-NH- or -NH-CO-NH-, the cyclisation of a compound of the formula I wherein R¹ is amino and R² is amino, (1-4C)alkylamino, (3-4C)alkenylamino, (3-4C)alkynylamino or a substituted-(1-4C)alkylamino with an appropriate carboxylic acid, an amide of a carboxylic acid, a urea or a carbonate.

The reaction is conveniently performed in the presence of a suitable inert solvent or diluent as defined hereinbefore and at a temperature in the range, for example, 10 to 150°C, preferably in the range 60 to 120°C.

(c) For the production of those optionally-substituted compounds of the formula I wherein R¹ and R² together form a group of the formula -N=N-NH-, the diazotisation and cyclisation of a compound of the formula I wherein R¹ is amino and R² is amino, (1-4C)alkylamino, (3-4C)alkenylamino, (3-4C)alkynylamino or a substituted-(1-4C)alkylamino.

A suitable diazotisation reagent is, for example, an alkali metal or alkaline earth metal nitrite, for example sodium nitrite, in the presence of a suitable acid such as sulphuric acid.

The reaction is conveniently performed at a temperature in the range, for example, -10 to +50°C, preferably in the range 0 to 30°C.

(d) For the production of those optionally-substituted compounds of the formula I wherein R¹ and R² together form a group of the formula -N=CH-CH=N-, the cyclisation of a compound of the formula I wherein R¹ is amino and R² is amino with an appropriate diketone.

The reaction is conveniently performed in the presence of a suitable inert solvent or diluent as described hereinbefore and at a temperature in the range, for example, 10 to 150°C, preferably in the range 50 to 100°C.

(e) For the production of those optionally-substituted compounds of the formula I wherein R¹ and R² together form a group of the formula -N=CH-CO-NH-, the cyclisation of a compound of the formula I wherein R¹ is amino and R² is amino, (1-4C)alkylamino, (3-4C)alkenylamino, (3-4C)alkynylamino or a substituted-(1-4C)alkylamino with an appropriate dicarboxylic acid or di-ester thereof.

The reaction is conveniently performed in the presence of a suitable inert solvent or diluent, as defined hereinbefore and at a temperature in the range, for example, 10 to 150°C, preferably in the range 50 to 100°C.

(f) For the production of those compounds of the formula I which bear a (1-4C)alkylsulphinyl or (1-4C)alkylsulphonyl substituent, the oxidation of a quinazoline derivative of the formula I which bears a (1-4C)alkylthio substituent.

[0027] A suitable oxidising agent is, for example, any agent known in the art for the oxidation of thio to sulphinyl and/or sulphonyl, for example, hydrogen peroxide, a peracid (such as 3-chloroperoxybenzoic or peroxyacetic acid), an alkali metal peroxysulphate (such as potassium peroxymonosulphate), chromium trioxide or gaseous oxygen in the presence of platinum. The oxidation is generally carried out under as mild conditions as possible and with the required stoichiometric amount of oxidising agent in order to reduce the risk of over oxidation and damage to other functional groups. In general the reaction is carried out in a suitable solvent or diluent such as methylene chloride, chloroform, acetone, tetrahydrofuran or *tert*-butyl methyl ether and at a temperature, for example, -25 to 50°C, conveniently at or near ambient temperature, that is in the range 15 to 35°C. When a compound carrying a sulphinyl group is required a milder oxidising agent may also be used, for example sodium or potassium metaperiodate, conveniently in a polar solvent such as acetic acid or ethanol. It will be appreciated that when a compound of the formula I containing a (1-4C)alkylsulphonyl group is required, it may be obtained by oxidation of the corresponding (1-4C)alkylsulphinyl compound as well as of the corresponding (1-4C)alkylthio compound.

[0028] When a pharmaceutically-acceptable salt of a tricyclic derivative of the formula I is required, for example an acid-addition salt of a tricyclic derivative of the formula I, it may be obtained, for example, by reaction of said compound with, for example, a suitable acid using a conventional procedure.

[0029] As stated hereinbefore the tricyclic derivative defined in the present invention possesses anti-cancer activity which is believed to arise from the receptor tyrosine kinase inhibitory activity of the compound. These properties may

be assessed, for example, using one or more of the procedures set out below:-

(a) An *in vitro* assay which determines the ability of a test compound to inhibit the enzyme receptor tyrosine kinase. Receptor tyrosine kinase was obtained in partially purified form from A-431 cells (derived from human vulval carcinoma) by procedures related to those described by Carpenter *et al.*, *J. Biol. Chem.*, 1979, 254, 4884, Cohen *et al.*, *J. Biol. Chem.*, 1982, 257, 1523 and by Braun *et al.*, *J. Biol. Chem.*, 1984, 259, 2051.

A-431 cells were grown to confluence using Dulbecco's modified Eagle's medium (DMEM) containing 5% fetal calf serum (FCS). The obtained cells were homogenised in a hypotonic borate/EDTA buffer at pH 10.1. The homogenate was centrifuged at 400 g for 10 minutes at 0-4°C. The supernatant was centrifuged at 25,000 g for 30 minutes at 0-4°C. The pelleted material was suspended in 30 mM Hepes buffer at pH 7.4 containing 5% glycerol, 4 mM benzamidine and 1% Triton X-100, stirred for 1 hour at 0-4°C, and recentrifuged at 100,000 g for 1 hour at 0-4°C. The supernatant, containing solubilised receptor tyrosine kinase, was stored in liquid nitrogen.

For test purposes 40 µl of the enzyme solution so obtained was added to a mixture of 400 µl of a mixture of 150 mM Hepes buffer at pH 7.4, 500 µM sodium orthovanadate, 0.1% Triton X-100, 10% glycerol, 200 µl water, 80 µl of 25 mM DTT and 80 µl of a mixture of 12.5 mM manganese chloride, 125 mM magnesium chloride and distilled water. There was thus obtained the test enzyme solution.

Each test compound was dissolved in dimethylsulphoxide (DMSO) to give a 50 mM solution which was diluted with 40 mM Hepes buffer containing 0.1% Triton X-100, 10% glycerol and 10% DMSO to give a 500 µM solution. Equal volumes of this solution and a solution of epidermal growth factor (EGF; 20 µg/ml) were mixed.

[γ-³²P]ATP (3000 Ci/mM, 250 µCi) was diluted to a volume of 2 ml by the addition of a solution of ATP (100 µM) in distilled water. An equal volume of a 4 mg/ml solution of the peptide Arg-Arg-Leu-Ile-Glu-Asp-Ala-Glu-Tyr-Ala-Ala-Arg-Gly in a mixture of 40 mM Hepes buffer at pH 7.4, 0.1% Triton X-100 and 10% glycerol was added.

The test compound/EGF mixture solution (5 µl) was added to the test enzyme solution (10 µl) and the mixture was incubated at 0-4°C for 30 minutes. The ATP/peptide mixture (10 µl) was added and the mixture was incubated at 25°C for 10 minutes. The phosphorylation reaction was terminated by the addition of 5% trichloroacetic acid (40 µl) and bovine serum albumin (BSA; 1 mg/ml, 5 µl). The mixture was allowed to stand at 4°C for 30 minutes and then centrifuged. An aliquot (40 µl) of the supernatant was placed onto a strip of Whatman p 81 phosphocellulose paper. The strip was washed in 75 mM phosphoric acid (4 x 10 ml) and blotted dry. Radioactivity present in the filter paper was measured using a liquid scintillation counter (Sequence A). The reaction sequence was repeated in the absence of the EGF (Sequence B) and again in the absence of the test compound (Sequence C).

Receptor tyrosine kinase inhibition was calculated as follows:-

$$\% \text{ Inhibition} = \frac{100 - (A-B)}{C - B} \times 100$$

The extent of inhibition was then determined at a range of concentrations of test compound to give an IC₅₀ value.

(b) An *in vitro* assay which determines the ability of a test compound to inhibit the growth of the human nasopharyngeal cancer cell line KB.

KB cells were seeded into wells at a density of 1×10^4 - 1.5×10^4 cells per well and grown for 24 hours in DMEM supplemented with 5% FCS (charcoal-stripped). Cell growth was determined after incubation for 3 days by the extent of metabolism of MTT tetrazolium dye to furnish a bluish colour. Cell growth was then determined in the presence of EGF (10 ng/ml) or in the presence of EGF (10 ng/ml) and a test compound at a range of concentrations. An IC₅₀ value could then be calculated.

(c) An *in vivo* assay in a group of male rats which determines the ability of a test compound (usually administered orally as a ball-milled suspension in 0.5% polysorbate) to inhibit the stimulation of liver hepatocyte growth caused by the administration of the growth factor TGFα (400 µg/kg subcutaneously, usually dosed twice, 3 and 7 hours respectively after the administration of the test compound).

[0030] In a control group of rats, the administration of TGFα causes on average a 5-fold stimulation of liver hepatocyte growth.

[0031] Cell-growth in the control and test animals is determined as follows:-

[0032] On the morning of the day after the dosing of the test compound (or 0.5% polysorbate in the control group), the animals are dosed with bromodeoxyuridine (BrdU; 100 mg/kg intraperitoneally). The animals are killed four hours later and the livers are excised. Slices are cut from each liver and the uptake of BrdU is determined by a conventional immunohistochemical technique similar to that described on pages 267 and 268 of an article by Goldsworthy *et al.* in *Chemically Induced Cell Proliferation: Implications for Risk Assessment*, Wiley-Liss Inc., 1991, pages 253-284. Further tests were carried out using a range of doses of the test compounds to allow the calculation of an approximate ED₅₀.

value for the inhibition of liver hepatocyte proliferation as determined by inhibition of the uptake of BrdU.

[0033] Although the pharmacological properties of the compounds of the formula I vary with structural change as expected, in general activity possessed by compounds of the formula I may be demonstrated at the following concentrations or doses in one or more of the above tests (a), (b) and (c):-

Test (a):- IC_{50} in the range, for example, 0.0005-1 μM ;

Test (b):- IC_{50} in the range, for example, 0.01-10 μM ;

Test (c):- ED_{50} in the range, for example, 1-100 mg/kg.

[0034] Thus, by way of example, the compound 3-methyl-8-(3'-methylanilino)-3H-imidazo[4,5-g]quinazoline has an IC_{50} of 0.035 μM in Test (a), an IC_{50} of 0.97 μM in Test (b) and an ED_{50} of <5 mg/kg in Test (c); and the compound 3-methyl-8-(3'-methylanilino)-1,2-dihydro-3H-imidazo[4,5-g]quinazolin-2-one has an IC_{50} of 0.016 μM in Test (a), an IC_{50} of 1.19 μM in Test (b) and an ED_{50} of <12.5 mg/kg in Test (c).

[0035] According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a tricyclic derivative of the formula I, or a pharmaceutically-acceptable salt thereof, as defined hereinbefore in association with a pharmaceutically-acceptable diluent or carrier.

[0036] The composition may be in a form suitable for oral administration, for example as a tablet or capsule, for parenteral injection (including intravenous, subcutaneous, intramuscular, intravascular or infusion) as a sterile solution, suspension or emulsion, for topical administration as an ointment or cream or for rectal administration as a suppository.

[0037] In general the above compositions may be prepared in a conventional manner using conventional excipients.

[0038] The compound will normally be administered to a warm-blooded animal at a unit dose within the range 5-5000 mg per square meter body area of the animal, i.e. approximately 0.1-100 mg/kg, and this normally provides a therapeutically-effective dose. A unit dose form such as a tablet or capsule will usually contain, for example 1-250 mg of active ingredient. Preferably a daily dose in the range of 1-50 mg/kg is employed. However the daily dose will necessarily be varied depending upon the host treated, the particular route of administration, and the severity of the illness being treated. Accordingly the optimum dosage may be determined by the practitioner who is treating any particular patient.

[0039] According to a further aspect of the present invention there is provided a tricyclic derivative of the formula I as defined hereinbefore for use in a method of treatment of the human or animal body by therapy.

[0040] We have now found that the compounds of the present invention possess anti-cancer properties which are believed to arise from their receptor tyrosine kinase inhibitory activity. Accordingly the compounds of the present invention are expected to be useful in the treatment of diseases or medical conditions mediated alone or in part by the enzyme receptor tyrosine kinase, i.e. the compounds may be used to produce a receptor tyrosine kinase inhibitory effect in a warm-blooded animal in need of such treatment. Thus the compounds of the present invention provide a method for treating the proliferation of malignant cells characterised by inhibition of the enzyme receptor tyrosine kinase, i.e. the compounds may be used to produce an anti-proliferative effect mediated alone or in part by the inhibition of the enzyme receptor tyrosine kinase. Accordingly the compounds of the present invention are expected to be useful in the treatment of cancer by providing an anti-proliferative effect, particularly in the treatment of receptor tyrosine kinase sensitive cancers such as cancers of the breast, lung, colon, rectum, stomach, prostate, bladder, pancreas and ovary.

[0041] Thus according to this aspect of the invention there is provided the use of a tricyclic derivative of the formula I, or a pharmaceutically-acceptable salt thereof, as defined hereinbefore in the manufacture of a medicament for use in the production of an anti-cancer effect in a warm-blooded animal such as man.

[0042] According to a further feature of this aspect of the invention there is provided a method for producing an anti-cancer effect in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a tricyclic derivative as defined immediately above.

[0043] As stated above the size of the dose required for the therapeutic or prophylactic treatment of a particular cancer will necessarily be varied depending on the host treated, the route of administration and the severity of the illness being treated. A unit dose in the range, for example, 1-100 mg/kg, preferably 1-50 mg/kg is envisaged.

[0044] The anti-cancer treatment defined hereinbefore may be applied as a sole therapy or may involve, in addition to the quinazoline derivative of the invention, one or more other anti-tumour substances, for example those selected from, for example, mitotic inhibitors, for example vinblastine; alkylating agents, for example cis-platin, carboplatin and cyclophosphamide; antimetabolites, for example 5-fluorouracil, cytosine arabinoside and hydroxyurea, or, for example, one of the preferred antimetabolites disclosed in European Patent Application No. 239362 such as N-[5-[N-(3,4-dihydro-2-methyl-4-oxoquinazolin-6-ylmethyl)-N-methylamino]-2-thenoyl]-L-glutamic acid; intercalating antibiotics, for example adriamycin and bleomycin; enzymes, for example asparaginase; topoisomerase inhibitors, for example etoposide; biological response modifiers, for example interferon; and anti-hormones, for example antiestrogens such as 'NOLVADEX' (tamoxifen) or, for example antiandrogens such as 'CASODEX' (4'-cyano-3-(4-fluorophenylsulphonyl)-2-hydroxy-2-methyl-3-(trifluoromethyl)propionanilide. Such conjoint treatment may be achieved by way of the simul-

taneous, sequential or separate dosing of the individual components of the treatment. According to this aspect of the invention there is provided a pharmaceutical product comprising a tricyclic derivative of the formula I as defined hereinbefore and an additional anti-tumour substance as defined hereinbefore for the conjoint treatment of cancer.

[0045] As stated above the tricyclic derivative defined in the present invention is an effective anti-cancer agent, which property is believed to arise from its receptor tyrosine kinase inhibitory properties. Such a tricyclic derivative of the invention is expected to possess a wide range of anti-cancer properties as receptor tyrosine kinases have been implicated in many common human cancers such as leukaemia and breast, lung, colon, rectal, stomach, prostate, bladder, pancreas and ovarian cancer. Thus it is expected that a tricyclic derivative of the invention will possess anti-cancer activity against these cancers. It is in addition expected that a tricyclic derivative of the present invention will possess activity against a range of leukaemias, lymphoid malignancies and solid tumours such as carcinomas and sarcomas in tissues such as the liver, kidney, prostate and pancreas.

[0046] The invention will now be illustrated in the following non-limiting Examples in which, unless otherwise stated:-

- (i) evaporations were carried out by rotary evaporation in vacuo and work-up procedures were carried out after removal of residual solids such as drying agents by filtration;
- (ii) operations were carried out at ambient temperature, that is in the range 18-25°C and under an atmosphere of an inert gas such as argon;
- (iii) column chromatography (by the flash procedure) and medium pressure liquid chromatography (HPLC) were performed on Merck Kieselgel silica (Art. 9385) or Merck Lichroprep RP-18 (Art. 9303) reversed-phase silica obtained from E. Merck, Darmstadt, Germany;
- (iv) yields are given for illustration only and are not necessarily the maximum attainable;
- (v) melting points are uncorrected and were determined using a Mettler SP62 automatic melting point apparatus, an oil-bath apparatus or a Koffler hot plate apparatus.
- (vi) the structures of the end-products of the formula I were confirmed by nuclear (generally proton) magnetic resonance (NMR) and mass spectral techniques; proton magnetic resonance chemical shift values were measured on the delta scale and peak multiplicities are shown as follows: s, singlet; d, doublet; t, triplet; m, multiplet;
- (vii) intermediates were not generally fully characterised and purity was assessed by thin layer chromatography (TLC), infra-red (IR) or NMR analysis;
- (viii) the following abbreviations have been used:-

DMF N,N-dimethylformamide;

DMA N,N-dimethylacetamide.

Example 1

[0047] A mixture of 6-amino-7-methylamino-4-(3'-methylanilino)quinazoline (0.61 g) and formic acid (50 ml) was stirred and heated to reflux for 1 hour. The mixture was evaporated and the residue was triturated under a dilute aqueous ammonium hydroxide solution. The resultant solid was isolated, washed with water and dried. There was thus obtained 3-methyl-8-(3'-methylanilino)-3H-imidazo[4,5-g]quinazoline (0.59 g), m.p. >290°C;

NMR Spectrum: (CD₃SOCD₃) 2.36 (s, 3H), 3.95 (s, 3H), 6.94 (d, 1H), 7.28 (m, 1H), 7.75 (m, 2H), 7.93 (s, 1H), 8.50 (s, 1H), 8.55 (s, 1H), 9.06 (s, 1H), 9.72 (broad s, 1H);

Elemental Analysis: Found C, 66.7; H, 5.4; N, 22.9;

C₁₇H₁₆N₆ 1H₂O requires C, 66.4; H, 5.5; N, 22.8%.

[0048] The 6-amino-7-methylamino-4-(3'-methylanilino)quinazoline used as a starting material was obtained as follows:

[0049] A mixture of 4-chloroanthranilic acid (17.2 g) and formamide (10 ml) was stirred and heated to 130°C for 45 minutes and to 175°C for 75 minutes. The mixture was allowed to cool to approximately 100°C and 2-(2-ethoxyethoxy) ethanol (50 ml) was added. The solution so formed was poured into a mixture (250 ml) of ice and water. The precipitate was isolated, washed with water and dried. There was thus obtained 7-chloroquinazolin-4-one (15.3 g, 85%).

[0050] A portion (6 g) of the material so obtained was added portionwise to a stirred mixture of concentrated sulphuric acid (12 ml) and fuming nitric acid (12 ml). The mixture was heated to 110°C for 30 minutes. The mixture was cooled to ambient temperature and poured onto ice. The solid was isolated, washed with water and dried. There was thus obtained 7-chloro-6-nitroquinazolin-4-one (6.89 g, 92%).

[0051] A mixture of a portion (4 g) of the material so obtained, thionyl chloride (30 ml), phosphoryl chloride (5 ml) and DMF (10 drops) was stirred and heated to reflux for 4 hours. The mixture was evaporated. A mixture of the residue, 3'-methylaniline (1.89 g) and isopropanol (25 ml) was stirred and heated to reflux for 2 hours. The mixture was filtered and the solid was washed with isopropanol and with diethyl ether. There was thus obtained 7-chloro-4-(3'-methylanilino)-6-nitroquinazoline (3.74 g, 67%), m.p. 271-274°C.

NMR Spectrum: (CD₃SOCD₃) 2.37 (s, 3H), 7.13 (d, 1H), 7.47 (t, 1H), 7.57 (m, 2H), 8.20 (s, 1H), 8.83 (s, 1H), 9.72 (s, 1H).

[0052] After repetition of the previous steps, a mixture of 7-chloro-4-(3'-methylanilino)-6-nitroquinazoline (10.5 g), an ethanolic solution of methylamine (30% weight/volume; 100 ml) and ethanol (100 ml) was stirred at ambient temperature for 16 hours. The mixture was evaporated to give 7-methylamino-4-(3'-methylanilino)-6-nitroquinazoline which

was used without further purification.
[0053] A mixture of 7-methylamino-4-(3'-methylanilino)-6-nitroquinazoline (0.7 g), 10% palladium-on-charcoal catalyst (0.07 g) and ethanol (100 ml) was stirred and heated to 45°C under an atmosphere of hydrogen for 2 hours. The mixture was cooled to ambient temperature and filtered. The filtrate was evaporated and the residue was purified by column chromatography using increasingly polar mixtures of methylene chloride and methanol as eluent. There was thus obtained 6-amino-7-methylamino-4-(3'-methylanilino)quinazoline (0.71 g) as a gum.

Example 2

[0054] A mixture of 6-amino-7-methylamino-4-(3'-methylanilino)quinazoline (0.2 g), urea (0.266 g) and DHA (8 ml) was stirred and heated to reflux for 2 hours. The mixture was evaporated and the residue was triturated under water. The resultant solid was isolated, washed with water and dried. The residue was dissolved in a mixture of DMSO (1 ml) and methanol (2 ml). The solution was acidified to pH1 by the addition of trifluoroacetic acid. The precipitate was isolated, washed with water, with acetone and with diethyl ether and dried. There was thus obtained 3-methyl-8-(3'-methylanilino)-1,2-dihydro-3H-imidazo[4,5-g]quinazolin-2-one (0.128 g);

NMR Spectrum: (CD₃SOCD₃) 3.38 (s, 3H), 3.40 (s, 3H), 7.11 (d, 1H), 7.36 (m, 1H), 7.39 (s, 1H), 7.50 (d, 1H), 7.52 (s, 1H), 8.24 (s, 1H), 8.79 (s, 1H), 10.9 (broad s, 1H), 11.95 (broad s, 1H);

Elemental Analysis: Found C, 54.6; H, 3.8; N, 16.7;

C₁₇H₁₅N₅O 1CF₃CO₂H requires C, 54.4; H, 3.8; N, 16.7%.

Example 3

[0055] A solution of sodium nitrite (0.065 g) in water (1 ml) was added dropwise during 5 minutes to a stirred suspension of 6-amino-4-(3'-chloro-4'-fluoroanilino)-7-methylaminoquinazoline (0.3 g) in 2N aqueous sulphuric acid solution (10 ml) which had been cooled to 0°C. The mixture was stirred at 0°C for 5 minutes, allowed to warm to ambient temperature and stirred for 20 minutes. The mixture was basified to pH10 by the addition of a concentrated aqueous ammonium hydroxide solution. The precipitate was isolated and triturated under methanol (10 ml). The solid was isolated, washed with methanol and with diethyl ether and dried. There was thus obtained 8-(3'-chloro-4'-fluoroanilino)-3-methyl-3H-[1,2,3]triazolo[4,5-g]quinazoline (0.177 g);

NMR Spectrum: (CD₃SOCD₃) 4.43 (s, 3H), 7.47 (m, 1H), 7.92 (m, 1H), 8.24 (s, 1H), 8.31 (m, 1H), 8.65 (s, 1H), 9.48 (s, 1H), 10.26 (broad s, 1H);

Elemental Analysis: Found C, 54.2; H, 3.0; N, 24.4;

C₁₆H₁₀N₆ClF 0.45CH₃OH requires C, 54.0; H, 3.4; N, 24.5%.

[0056] The 6-amino-4-(3'-chloro-4'-fluoroanilino)-7-methylaminoquinazoline used as a starting material was obtained as follows:-

[0057] A mixture of 7-chloro-6-nitroquinazolin-4-one (30 g), thionyl chloride (300 ml) and DMF (0.5 ml) was stirred and heated to reflux for 5 hours. The mixture was evaporated, toluene (50 ml) was added and the solution was evaporated. A mixture of the residue, 3-chloro-4-fluoroaniline (19.5 g) and isopropanol (100 ml) was stirred and heated to reflux for 2 hours. The mixture was cooled to ambient temperature. The precipitate was isolated and washed with isopropanol and with diethyl ether. There was thus obtained 7-chloro-4-(3'-chloro-4'-fluoroanilino)-6-nitroquinazoline hydrochloride (23.2 g);

NMR Spectrum: (CD₃SOCD₃) 7.50 (m, 1H), 7.82 (m, 1H), 8.14 (m, 1H), 8.18 (s, 1H), 8.88 (s, 1H), 9.67 (s, 1H), 11.3 (broad s, 1H);

Elemental Analysis: Found C, 43.9; H, 2.1; N, 14.5;

C₁₄H₈N₄Cl₂F 0.8HCl requires C, 43.8; H, 2.3; N, 14.6%.

[0058] A mixture of a portion (10 g) of the quinazoline so obtained, an ethanolic solution of methylamine (33% weight/volume, 50 ml) and ethanol (100 ml) was stirred and heated to 70°C for 5 hours under a condenser cooled with solid carbon dioxide. The mixture was allowed to stand at ambient temperature for 16 hours. The precipitate was isolated, washed with ethanol and with diethyl ether and dried. There was thus obtained 4-(3'-chloro-4'-fluoroanilino)-7-methylamino-6-nitroquinazoline (5.8 g);

NMR Spectrum: (CD₃SOCD₃) 2.99 (d, 3H), 6.89 (s, 1H), 7.43 (m, 1H), 7.81 (m, 1H), 7.96 (m, 1H), 8.16 (m, 1H), 8.49 (s, 1H), 9.45 (s, 1H), 10.2 (broad s, 1H).

[0059] A mixture of a portion (4.4 g) of the quinazoline so obtained, 10% palladium-on-charcoal catalyst (0.6 g), methylene chloride (150 ml) and ethanol (150 ml) was stirred under an atmosphere of hydrogen for 5 hours. The mixture

was filtered and the filtrate was evaporated. The residue was dissolved in a mixture of methylene chloride (50 ml) and methanol (10 ml) and the solution was stood at ambient temperature for 3 hours. A precipitate was deposited. The mixture was filtered and the filtrate was evaporated. The residue from the filtrate was purified by column chromatography using a 19:1 mixture of methylene chloride and methanol as eluent. There was thus obtained 6-amino-4-(3'-chloro-4'-fluoroanilino)-7-methylaminoquinazoline (1.84 g), m.p. 244-247°C;

NMR Spectrum (CD_3SOCD_3) 2.87 (d, 3H), 5.16 (broad s, 2H), 5.97 (m, 1H), 6.55 (s, 1H), 7.28 (m, 2H), 7.36 (m, 1H), 8.16 (m, 1H), 8.31 (s, 1H), 9.25 (broad s, 1H);

Elemental Analysis: Found C, 53.7; H, 5.0; N, 19.0;

$\text{C}_{15}\text{H}_{13}\text{N}_5\text{ClF}$ 1.5 CH_3OH requires C, 54.1; H, 5.2; N, 19.1%.

Example 4

[0060] A mixture of 6-amino-4-(3'-chloro-4'-fluoroanilino)-7-methylaminoquinazoline (0.3 g) and trifluoroacetic acid (10 ml) was stirred and heated to reflux for 30 minutes. The mixture was cooled to ambient temperature and evaporated. The residue was triturated under dilute aqueous ammonium hydroxide solution. The residue was washed with water and dried. There was thus obtained 8-(3'-chloro-4'-fluoroanilino)-3-methyl-2-trifluoromethyl-3H-imidazo[4,5-g]quinazoline (0.182 g);

NMR Spectrum: (CD_3SOCD_3) 4.08 (s, 3H), 7.46 (t, 1H), 7.92 (m, 1H), 8.20 (s, 1H), 8.28 (m, 1H), 8.64 (s, 1H), 9.19 (s, 1H), 10.04 (broad s, 1H); Elemental Analysis: Found C, 49.5; H, 2.4; N, 16.8;

$\text{C}_{17}\text{H}_{10}\text{N}_5\text{ClF}_4$ 0.2 $\text{CF}_3\text{CO}_2\text{H}$ requires C, 49.9; H, 2.4; N, 16.7%.

Example 5

[0061] A mixture of 6-amino-4-(3'-chloro-4'-fluoroanilino)-7-methylaminoquinazoline (0.3 g), tetraethyl orthocarbonate (2 ml) and acetic acid (0.07 ml) was stirred and heated to 100°C for 2 hours. The mixture was cooled to ambient temperature. The precipitate was isolated, washed with diethyl ether and dried. There was thus obtained 8-(3'-chloro-4'-fluoroanilino)-2-ethoxy-3-methyl-3H-imidazo[4,5-g]quinazoline (0.253 g);

NMR Spectrum: (CD_3SOCD_3) 1.47 (t, 3H), 3.64 (s, 3H), 4.65 (m, 2H), 7.37 (s, 1H), 7.43 (m, 1H), 7.92 (m, 1H), 8.32 (m, 1H), 8.56 (s, 1H), 8.60 (s, 1H), 9.67 (broad s, 1H);

Elemental Analysis: Found C, 58.3; H, 4.0; N, 18.9;

$\text{C}_{18}\text{H}_{15}\text{N}_5\text{ClFO}$ requires C, 58.1; H, 4.1; N, 18.8%.

Example 6

[0062] A mixture of 6-amino-7-(3-dimethylaminopropylamino)-4-(3'-methylanilino)quinazoline (0.15 g) and formic acid (2 ml) was stirred and heated to reflux for 2 hours. The mixture was evaporated and the residue was triturated under diethyl ether. The resultant solid was filtered off and washed with diethyl ether. There was thus obtained 3-(3-dimethylaminopropyl)-8-(3'-methylanilino)-3H-imidazo[4,5-g]quinazoline as a solid (0.14 g);

NMR Spectrum: (CD_3SOCD_3) 2.02 (m, 2H), 2.19 (s, 6H), 2.25 (t, 2H), 2.34 (s, 3H), 4.39 (t, 2H), 6.94 (d, 1H), 7.28 (m, 1H), 7.75 (m, 2H), 7.97 (s, 1H), 8.55 (s, 2H), 9.02 (s, 1H), 9.68 (broad s, 1H).

[0063] The 6-amino-7-(3-dimethylaminopropylamino)-4-(3'-methylanilino)quinazoline used as a starting material was obtained as follows:-

[0064] A mixture of 7-chloro-4-(3'-methylanilino)-6-nitroquinazoline (1.2 g), 3-dimethylaminopropylamine (6 ml) and DHA (20 ml) was stirred and heated to 80°C for 2 hours. The mixture was evaporated and the residue was triturated under water. The resultant solid was washed with water and dried. There was thus obtained 7-(3-dimethylaminopropylamino)-6-nitro-4-(3'-methylanilino)quinazoline (1.35 g) which was used without further purification.

[0065] A mixture of a portion (1 g) of the material so obtained, 10% palladium-on-charcoal catalyst (0.1 g) and ethanol (80 ml) was stirred and heated to 45°C for 2 hours. The mixture was cooled to ambient temperature and filtered. The filtrate was evaporated. The residue was partitioned between ethyl acetate and 1N aqueous hydrochloric acid solution. The aqueous layer was basified by the addition of ammonium hydroxide solution and the mixture was extracted with ethyl acetate. The organic phase was dried (MgSO_4) and evaporated. There was thus obtained 6-amino-7-(3-dimethylaminopropylamino)-4-(3'-methylanilino)-quinazoline as a foam (0.61 g), m.p. 60-66°C.

Example 7

[0066] A mixture of 6,7-diamino-4-(3'-chloro-4'-fluoroanilino)quinazoline (0.1 g), biacetyl (0.037 g) and ethanol (2 ml) was heated to reflux for 4 hours. The mixture was cooled to ambient temperature and the precipitate was isolated, washed with water and dried. There was thus obtained 4-(3'-chloro-4'-fluoroanilino)-7,8-dimethylpyrazino-[2,3-g]quina-

zoline (0.054 g), m.p. >270°C;

NMR Spectrum: (CD₃SOCD₃) 2.77 (s, 6H), 7.49 (m, 1H), 7.92 (m, 1H), 8.26 (s, 1H), 8.35 (d, 1H), 8.70 (s, 1H), 9.37 (s, 1H), 10.27 (broad s, 1H).

[0067] The 6,7-diamino-4-(3'-chloro-4'-fluoroanilino)quinazoline used as a starting material was obtained as follows:-

[0068] Sodium azide (0.509 g) was added portionwise to a stirred solution of 7-chloro-4-(3'-chloro-4'-fluoroanilino)-6-nitroquinazoline hydrochloride (1 g) in DMA (250 ml). The mixture was stirred and heated to 90°C for 2 hours. A quantity (0.2 g) of 10% palladium-on-charcoal catalyst was added and the mixture was stirred under an atmosphere of hydrogen and heated to 80°C for 2 hours. The mixture was cooled to ambient temperature and filtered. The filtrate was evaporated and the residue was purified by column chromatography using a 19:1 mixture of methylene chloride and methanol as eluent. There was thus obtained 6,7-diamino-4-(3'-chloro-4'-fluoroanilino)-quinazoline as a solid (0.39 g), m.p. 254-257°C;




NMR Spectrum: (CD₃SOCD₃) 5.17 (broad s, 2H), 5.83 (broad s, 1H), 6.78 (s, 1H), 7.28 (s, 1H), 7.3 (broad s, 1H), 7.36 (m, 1H), 7.76 (m, 1H), 8.15 (m, 1H), 8.28 (s, 1H).

Example 8

[0069] A mixture of 6-amino-4-(3'-chloro-4'-fluoroanilino)-7-methylaminoquinazoline (0.05 g), oxalic acid (0.075 g), water (1 ml) and acetic acid (1 ml) was stirred and heated to reflux for 3 hours. The mixture was evaporated and the residue was triturated with water. The resultant solid was washed with water and dried. There was thus obtained 4-(3'-chloro-4'-fluoroanilino)-7-hydroxy-9-methyl-8-oxo-8,9-dihydropyrazino[2,3-g]quinazoline (0.028 g);


NMR Spectrum: (CD₃SOCD₃) 3.62 (s, 3H), 7.43 (t, 1H), 7.63 (s, 1H), 7.75 (m, 1H), 7.99 (s, 1H), 8.10 (m, 1H), 8.59 (s, 1H), 10.0 (broad s, 1H), 11.9 (broad s, 1H).

CHEMICAL FORMULAE



Claims

1. A tricyclic derivative of the formula I



I

wherein R^1 and R^2 together form a group of the formula $-N=CH-NH-$, $-N=CH-O-$, $-N=CH-S-$, $-N=N-NH-$, $-NH-N=CH-$, $-NH-CH=CH-$, $-NH-CO-NH-$, $-NH-CO-O-$, $-NH-CO-S-$, $-NH-NH-CO-$, $-N=CH-CH=CH-$, $-N=N-CH=CH-$, $-N=CH-N=CH-$, $-N=CH-CH=N-$, $-NH-CO-CH=CH-$ or $-N=CH-CO-NH-$ (with in each case a nitrogen atom being located at the 6-position of the quinazoline ring) and the 5- or 6-membered ring so formed may optionally bear one or two substituents, any substituent on an available nitrogen atom being selected from (1-4C)alkyl, (3-4C)alkenyl, (3-4C)alkynyl, halogeno-(1-4C)alkyl, hydroxy-(1-4C)alkyl, (2-4C)alkanoyloxy-(1-4C)alkyl, (1-4C)alkoxy-(1-4C)alkyl, cyano-(1-4C)alkyl, amino-(1-4C)alkyl, (1-4C)alkylamino-(1-4C)alkyl and di-[(1-4C)alkyl]amino-(1-4C)alkyl, and any substituent on an available carbon atom being selected from halogeno, amino, hydroxy, carbamoyl, cyano, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, (1-4C)alkoxy, (1-4C)alkylthio, (1-4C)alkylsulphinyl, (1-4C)alkylsulphonyl, (1-4C)alkylamino, di-[(1-4C)alkyl]amino, (2-4C)alkanoyl, N -(1-4C)alkylcarbamoyl, N,N -di-[(1-4C)alkyl]carbamoyl, halogeno-(1-4C)alkyl, hydroxy-(1-4C)alkyl, (2-4C)alkanoyloxy-(1-4C)alkyl, (1-4C)alkoxy-(1-4C)alkyl, cyano-(1-4C)alkyl, amino-(1-4C)alkyl, (1-4C)alkylamino-(1-4C)alkyl and di-[(1-4C)alkyl]amino-(1-4C)alkyl; and m is the integer 1, 2 or 3 and each R^3 is independently hydrogen, halogeno, trifluoromethyl, hydroxy, amino, nitro, cyano, (1-4C)alkyl, (1-4C)alkoxy, (1-4C)alkylamino, di-[(1-4C)alkyl]amino or (2-4C)alkanoylamino; or a pharmaceutically-acceptable salt thereof.

2. A tricyclic derivative of the formula I as claimed in claim 1 wherein the 6,6,5-tricyclic ring defined by the linking of the groups R^1 and R^2 is selected from 3H-imidazo[4,5-g]quinazolin-8-yl, oxazolo[4,5-g]quinazolin-8-yl, thiazolo[4,5-g]quinazolin-8-yl, 3H-[1,2,3]triazolo[4,5-g]quinazolin-8-yl, 1H-pyrazolo[3,4-g]quinazolin-8-yl, 6H-pyrrolo[2,3-g]quinazolin-4-yl, 2-oxo-1,2-dihydro-3H-imidazo[4,5-g]quinazolin-8-yl, 2-oxo-1,2-dihydrooxazolo[4,5-g]quinazolin-8-yl, 2-oxo-1,2-dihydrothiazolo[4,5-g]quinazolin-8-yl and 3-oxo-2,3-dihydro-1H-pyrazolo[3,4-g]quinazolin-8-yl,


and the 5-membered ring involving R^1 and R^2 may optionally bear one or two substituents, any substituent on an available nitrogen atom being selected from methyl, ethyl, propyl, allyl, prop-2-ynyl, 2,2,2-trifluoroethyl, 2-hydroxyethyl, 3-hydroxypropyl, 2-acetoxyethyl, 2-methoxyethyl, 3-methoxypropyl, cyanomethyl, 2-cyanoethyl, 2-aminoethyl, 2-methylaminoethyl and 2-dimethylaminoethyl, and any substituent on an available carbon atom being selected from fluoro, chloro, amino, carbamoyl, cyano, methyl, ethyl, propyl, vinyl, allyl, ethynyl, prop-2-ynyl, methoxy, ethoxy, propoxy, methylthio, methylsulphinyl, methylsulphonyl, methylamino, dimethylamino, acetyl, N -methylcarbamoyl, N,N -dimethylcarbamoyl, fluoromethyl, difluoromethyl, trifluoromethyl, 2,2,2-trifluoroethyl, hydroxymethyl, 2-hydroxyethyl, acetoxyethyl, 2-acetoxyethyl, methoxymethyl, 2-methoxyethyl, cyanomethyl, 2-cyanoethyl, aminomethyl, 2-aminoethyl, methylaminomethyl, 2-methylaminoethyl, dimethylaminomethyl and 2-dimethylaminoethyl; and m is the integer 1, 2 or 3 and each R^3 is independently hydrogen, fluoro, chloro, bromo, trifluoromethyl, hydroxy, amino, nitro, cyano, methyl, ethyl, methoxy, methylamino, dimethylamino or acetamido; or a pharmaceutically-acceptable salt thereof.

3. A tricyclic derivative of the formula I as claimed in claim 1 wherein the 6,6,6-tricyclic ring defined by the linking of the groups R^1 and R^2 is selected from pyrido[2,3-g]quinazolin-4-yl, pyrimidino[4,5-g]cinnolin-9-yl, pyrimidino[4,5-g]quinazolin-4-yl, pyrazino[2,3-g]quinazolin-4-yl, 7-oxo-6,7-dihydropyrido[2,3-g]quinazolin-4-yl and 8-oxo-8,9-dihydropyrazino[2,3-g]quinazolin-4-yl, and the 6-membered ring involving R^1 and R^2 may optionally bear one or two substituents, any substituent on an available nitrogen atom being selected from methyl, ethyl and propyl, and any substituent on an available carbon atom being selected from fluoro, chloro, hydroxy, carbamoyl, cyano, methyl, methoxy, ethoxy, N -methylcarbamoyl,


N,N-dimethylcarbamoyl, trifluoromethyl and 2,2,2-trifluoroethyl; and m is the integer 1 or 2 and each R^3 is independently hydrogen, fluoro, chloro, bromo, trifluoromethyl, nitro, cyano, methyl or ethyl; or a pharmaceutically-acceptable salt thereof.

4. A tricyclic derivative of the formula I as claimed in claim 1 wherein the 6,6,5-tricyclic ring defined by the linking of the groups R^1 and R^2 is selected from 3-methyl-3H-imidazo[4,5-g]quinazolin-8-yl, 3-methyl-3H-[1,2,3]triazolo[4,5-g]quinazolin-8-yl and 3-methyl-2-oxo-1,2-dihydro-3H-imidazo[4,5-g]quinazolin-8-yl; and $(R^3)_m$ is 3'-methyl, 3'-chloro or 3'-chloro-4'-fluoro; or a pharmaceutically-acceptable salt thereof.
5. A tricyclic derivative of the formula I as claimed in claim 1 wherein the 6,6,5-tricyclic ring defined by the linking of the groups R^1 and R^2 is selected from 3-methyl-3H-imidazo[4,5-g]quinazolin-8-yl, 3-methyl-2-oxo-1,2-dihydro-3H-imidazo[4,5-g]quinazolin-8-yl and 3-methyl-2-trifluoromethyl-3H-imidazo[4,5-g]quinazolin-8-yl; and $(R^3)_m$ is 3'-methyl, 3'-chloro or 3'-chloro-4'-fluoro; or a pharmaceutically-acceptable salt thereof.
6. A tricyclic derivative of the formula I as claimed in claim 1 wherein the 6,6,6-tricyclic ring defined by the linking of the groups R^1 and R^2 is selected from 7,8-dimethylpyrazino[2,3-g]quinazolin-4-yl and 7-hydroxy-9-methyl-8-oxo-8,9-dihydropyrazino[2,3-g]quinazolin-4-yl; and $(R^3)_m$ is 3'-methyl, 3'-chloro or 3'-chloro-4'-fluoro; or a pharmaceutically-acceptable salt thereof.
7. A tricyclic derivative of the formula I as claimed in claim 1 selected from:-
3-methyl-8-(3'-methylanilino)-3H-imidazo[4,5-g]quinazoline,
3-methyl-8-(3'-methylanilino)-1,2-dihydro-3H-imidazo[4,5-g]quinazolin-2-one
and 8-(3'-chloro-4'-fluoroanilino)-3-methyl-2-trifluoromethyl-3H-imidazo[4,5-g]quinazoline;
or a pharmaceutically-acceptable salt thereof.
8. A tricyclic derivative of the formula I as claimed in claim 1 selected from:-
4-(3'-chloro-4'-fluoroanilino)-7,8-dimethylpyrazino[2,3-g]quinazoline and
4-(3'-chloro-4'-fluoroanilino)-7-hydroxy-9-methyl-8-oxo-8,9-dihydropyrazino[2,3-g]quinazoline;
or a pharmaceutically-acceptable salt thereof.
9. A process for the preparation of a tricyclic derivative of the formula I, or a pharmaceutically-acceptable salt thereof, as claimed in claim 1 which comprises:-

(a) the reaction of a quinazoline of the formula II



wherein Z is a displaceable group, with an aniline of the formula III



(b) for the production of those optionally-substituted compounds of the formula I wherein R^1 and R^2 together form a group of the formula $-N=CH-NH-$ or $-NH-CO-NH-$, the cyclisation of a compound of the formula I wherein R^1 is amino and R^2 is amino, (1-4C)alkylamino, (3-4C)alkenylamino, (3-4C)alkynylamino or a substituted-(1-4C)alkylamino with an appropriate carboxylic acid, an amide of a carboxylic acid, a urea or a carbonate;

(c) for the production of those optionally-substituted compounds of the formula I wherein R^1 and R^2 together form a group of the formula $-N=N-NH-$, the diazotisation and cyclisation of a compound of the formula I wherein R^1 is amino and R^2 is amino, (1-4C)alkylamino, (3-4C)alkenylamino, (3-4C)alkynylamino or a substituted-(1-4C)alkylamino;

(d) for the production of those optionally-substituted compounds of the formula I wherein R^1 and R^2 together form a group of the formula $-N=CH-CH=N-$, the cyclisation of a compound of the formula I wherein R^1 is amino and R^2 is amino with an appropriate diketone;

(e) for the production of those optionally-substituted compounds of the formula I wherein R^1 and R^2 together form a group of the formula $-N=CH-CO-NH-$, the cyclisation of a compound of the formula I wherein R^1 is amino and R^2 is amino, (1-4C)alkylamino, (3-4C)alkenylamino, (3-4C)alkynylamino or a substituted-(1-4C)alkylamino with an appropriate dicarboxylic acid or di-ester thereof; or

(f) for the production of those compounds of the formula I which bear a (1-4C)alkylsulphonyl or (1-4C)alkylsulphonyl substituent, the oxidation of a quinazoline derivative of the formula I which bears a (1-4C)alkylthio substituent;


and when a pharmaceutically-acceptable salt of a tricyclic derivative of the formula I is required it may be obtained using a conventional procedure.

10. A pharmaceutical composition which comprises a tricyclic derivative of the formula I, or a pharmaceutically-acceptable salt thereof, as claimed in any one of claims 1 to 8 in association with a pharmaceutically-acceptable diluent or carrier.

11. The use of a tricyclic derivative of the formula I, or a pharmaceutically-acceptable salt thereof, as claimed in any one of claims 1 to 8 in the manufacture of a medicament for use in the production of an anti-cancer effect in a warm-blooded animal.

Patentansprüche

1. Tricyclisches Derivat der Formel I



wobei R^1 und R^2 zusammen einen Rest der Formel $-N=CH-NH-$, $-N=CH-O-$, $-N=CH-S-$, $-N=N-NH-$, $-NH-N=CH-$, $-NH-CH=CH-$, $-NH-CO-NH-$, $-NH-CO-O-$, $-NH-CO-S-$, $-NH-NH-CO-$, $-N=CH-CH=CH-$, $-N=N-CH=CH-$, $-N=CH-N=CH-$, $-N=CH-CH=N-$, $-NH-CO-CH=CH-$ oder $-N=CH-CO-NH-$ (wobei sich in jedem Fall ein Stickstoffatom an der 6-Stellung des Chinazolinrings befindet) bilden und der so erzeugte 5- oder 6-gliedrige Ring gegebenenfalls einen oder zwei Substituenten tragen kann, wobei jeder Substituent an einem verfügbaren Stickstoffatom aus (1-4C)Alkyl-, (3-4C)Alkenyl-, (3-4C)Alkynyl-, Halogen-(1-4C)alkyl-, Hydroxy-(1-4C)alkyl-, (2-4C)Alkanoyloxy-(1-4C)alkyl-, (1-4C)Alkoxy-(1-4C)alkyl-, Cyano-(1-4C)alkyl-, Amino-(1-4C)alkyl-, (1-4C)Alkylamino-(1-4C)alkyl- und Di-[(1-4C)alkyl]amino-(1-4C)alkylresten ausgewählt ist und jeder Substituent an einem verfügbaren Kohlenstoffatom aus Halogenatomen, Amino-, Hydroxygruppen, Carbamoylresten, Cyanogruppen, (1-4C)Alkyl-, (2-4C)Alkenyl-, (2-4C)Alkynyl-, (1-4C)Alkoxy-, (1-4C)Alkylthio-, (1-4C)Alkylsulfonyl-, (1-4C)Alkylsulfonyl-, (1-4C)Alkylamino-, Di-[(1-4C)alkyl]amino-, (2-4C)Alkanoyl-, N -(1-4C)Alkylcarbamoyl-, N,N -Di[(1-4C)alkyl]carbamoyl-, Halogen-(1-4C)alkyl-, Hydroxy-(1-4C)alkyl-, (2-4C)Alkanoyloxy-(1-4C)alkyl-, (1-4C)Alkoxy-(1-4C)alkyl-, Cyano-(1-4C)alkyl-, Amino-(1-4C)alkyl-, (1-4C)Alkylamino-(1-4C)alkyl- und Di-[(1-4C)alkyl]amino-(1-4C)alkylresten ausgewählt ist; und m die ganze Zahl 1, 2 oder 3 ist, und jeder Rest R^3 unabhängig ein Wasserstoff-, Halogenatom, eine Trifluormethyl-, Hydroxy-, Amino-, Nitro-, Cyanogruppe, ein (1-4C)Alkyl-, (1-4C)Alkoxy-, (1-4C)Alkylamino-, Di-[(1-4C)alkyl]amino- oder (2-4C)Alkanoylaminorest ist; oder ein pharmazeutisch verträgliches Salz davon.


2. Tricyclisches Derivat der Formel I, wie in Anspruch 1 beansprucht, wobei der 6,6,5-tricyclische Ring, definiert durch die Verknüpfung der Reste R^1 und R^2 , aus 3H-Imidazo[4,5-g]chinazolin-8-yl-, Oxazolo[4,5-g]chinazolin-8-yl-, Thiazolo[4,5-g]chinazolin-8-yl-, 3H-[1,2,3]Triazolo[4,5-g]chinazolin-8-yl-, 1H-Pyrazolo[3,4-g]chinazolin-8-yl-, 6H-Pyrazolo[2,3-g]chinazolin-4-yl-, 2-Oxo-1,2-dihydro-3H-imidazo[4,5-g]chinazolin-8-yl-, 2-Oxo-1,2-dihydrooxazolo[4,5-g]chinazolin-8-yl-, 2-Oxo-1,2-dihydrothiazolo[4,5-g]chinazolin-8-yl- und 3-Oxo-2,3-dihydro-1H-pyrazolo[3,4-g]chinazolin-8-ylresten ausgewählt ist, und der 5-gliedrige Ring, der R^1 und R^2 enthält, gegebenenfalls einen oder zwei Substituenten tragen kann, wobei jeder Substituent an einem verfügbaren Stickstoffatom aus Methyl-, Ethyl-, Propyl-, Allyl-, Prop-2-ynyl-, 2,2,2-Trifluorethyl-, 2-Hydroxyethyl-, 3-Hydroxypropyl-, 2-Acetoxyethyl-, 2-Methoxyethyl-, 3-Methoxypropyl-, Cyanomethyl-, 2-Cyanoethyl-, 2-Aminoethyl-, 2-Methylaminoethyl- und 2-Dimethylaminoethylgruppen ausgewählt ist, und jeder Substituent an einem verfügbaren Kohlenstoffatom aus Fluor-, Chloratomen, Amino-, Carbamoyl-, Cyano-, Methyl-, Ethyl-, Propyl-, Vinyl-, Allyl-, Ethinyl-, Prop-2-ynyl-, Methoxy-, Ethoxy-, Propoxy-, Methylthio-, Methylsulfinyl-, Methylsulfonyl-, Methylamino-, Dimethylamino-, Acetyl-, N -Methylcarbamoyl-, N,N -Dimethylcarbamoyl-, Fluormethyl-, Difluormethyl-, Trifluormethyl-, 2,2,2-Trifluorethyl-, Hydroxymethyl-, 2-Hydroxyethyl-, Acetoxymethyl-, 2-Acetoxyethyl-, Methoxymethyl-, 2-Methoxyethyl-, Cyanomethyl-, 2-Cyanoethyl-, Aminomethyl-, 2-Aminoethyl-, Methylaminomethyl-, 2-Methylaminoethyl-, Dimethylaminomethyl- und 2-Dimethylaminoethylgruppen ausgewählt ist; und m die ganze Zahl 1, 2 oder 3 bedeutet, und jeder Rest R^3 unabhängig ein Wasserstoff-, Fluor-, Chlor-, Bromatom, eine Trifluormethyl-, Hydroxy-, Amino-, Nitro-, Cyano-, Methyl-, Ethyl-, Methoxy-, Methylamino-, Dimethylamino- oder Acetamidogruppe ist; oder ein pharmazeutisch verträgliches Salz davon.

3. Tricyclisches Derivat der Formel I, wie in Anspruch 1 beansprucht, wobei der 6,6,6-tricyclische Ring, definiert durch die Verknüpfung der Reste R^1 und R^2 , aus Pyrido[2,3-g]chinazolin-4-yl-, Pyrimidino[4,5-g]cinnolin-9-yl-, Pyrimidino[4,5-g]chinazolin-4-yl-, Pyrazino[2,3-g]chinazolin-4-yl-, 7-Oxo-6,7-dihydropyrido[2,3-g]chinazolin-4-yl- und 8-Oxo-8,9-dihydropyrazino[2,3-g]chinazolin-4-ylresten ausgewählt ist, und der 6-gliedrige Ring, der R^1 und R^2 enthält, gegebenenfalls einen oder zwei Substituenten tragen kann, wobei jeder Substituent an einem verfügbaren Stickstoffatom aus Methyl-, Ethyl- und Propylgruppen ausgewählt ist, und jeder Substituent an einem verfügbaren Kohlenstoffatom aus Fluor-, Chloratomen, Hydroxy-, Carbamoyl-, Cyano-, Methyl-, Methoxy-, Ethoxy-, N -Methylcarbamoyl-, N,N -Dimethylcarbamoyl-, Trifluormethyl- und 2,2,2-Trifluorethylgruppen ausgewählt ist; und m die ganze Zahl 1 oder 2 bedeutet, und jeder Rest R^3 unabhängig ein Wasserstoff-, Fluor-, Chlor-, Bromatom, eine Trifluormethyl-, Nitro-, Cyano-, Methyl- oder Ethylgruppe ist; oder ein pharmazeutisch verträgliches Salz davon.


4. Tricyclisches Derivat der Formel I, wie in Anspruch 1 beansprucht, wobei der 6,6,5-tricyclische Ring, definiert durch die Verknüpfung der Reste R^1 und R^2 , aus 3-Methyl-3H-imidazo[4,5-g]chinazolin-8-yl-, 3-Methyl-3H-[1,2,3]Triazolo[4,5-g]chinazolin-8-yl- und 3-Methyl-2-oxo-1,2-dihydro-3H-imidazo[4,5-g]chinazolin-8-ylresten ausgewählt ist; und $(R^3)_m$ eine 3'-Methylgruppe, ein 3'-Chloratom oder ein 3'-Chlor-4'-fluorrest ist; oder ein pharmazeutisch verträgliches Salz davon.

5. Tricyclisches Derivat der Formel I, wie in Anspruch 1 beansprucht, wobei der 6,6,5-tricyclische Ring, definiert durch die Verknüpfung der Reste R^1 und R^2 , aus 3-Methyl-3H-imidazo[4,5-g]chinazolin-8-yl-, 3-Methyl-2-oxo-1,2-dihydro-3H-imidazo[4,5-g]chinazolin-8-yl- und 3-Methyl-2-trifluormethyl-3H-imidazo[4,5-g]chinazolin-8-ylresten ausgewählt ist; und $(R^3)_m$ eine 3'-Methylgruppe, ein 3'-Chloratom oder ein 3'-Chlor-4'-fluorrest ist; oder ein pharmazeutisch verträgliches Salz davon.
6. Tricyclisches Derivat der Formel I, wie in Anspruch 1 beansprucht, wobei der 6,6,6-tricyclische Ring, definiert durch die Verknüpfung der Reste R^1 und R^2 , aus 7,8-Dimethylpyrazino[2,3-g]chinazolin-4-yl- und 7-Hydroxy-9-methyl-8-oxo-8,9-dihydropyrazino[2,3-g]chinazolin-4-ylresten ausgewählt ist; und $(R^3)_m$ eine 3'-Methylgruppe, ein 3'-Chloratom oder ein 3'-Chlor-4'-fluorrest ist; oder ein pharmazeutisch verträgliches Salz davon.
7. Tricyclisches Derivat der Formel I, wie in Anspruch 1 beansprucht, ausgewählt aus: 3-Methyl-8-(3'-methylanilino)-3H-imidazo[4,5-g]chinazolin, 3-Methyl-8-(3'-methylanilino)-1,2-dihydro-3H-imidazo[4,5-g]chinazolin-2-on und 8-(3'-Chlor-4'-fluoranilino)-3-methyl-2-trifluormethyl-3H-imidazo[4,5-g]chinazolin; oder einem pharmazeutisch verträglichen Salz davon.
8. Tricyclisches Derivat der Formel I, wie in Anspruch 1 beansprucht, ausgewählt aus: 4-(3'-Chlor-4'-fluoranilino)-7,8-dimethylpyrazino[2,3-g]chinazolin und 4-(3'-Chlor-4'-fluoranilino)-7-hydroxy-9-methyl-8-oxo-8,9-dihydropyrazino[2,3-g]chinazolin; oder einem pharmazeutisch verträglichen Salz davon.
9. Verfahren zur Herstellung eines tricyclischen Derivats der Formel I oder eines pharmazeutisch verträglichen Salzes davon, wie in Anspruch 1 beansprucht, das umfaßt:

(a) die Reaktion eines Chinazolins der Formel II



wobei Z eine Abgangsgruppe ist, mit einem Anilin der Formel III



- (b) zur Herstellung jener gegebenenfalls substituierten Verbindungen der Formel I, wobei R^1 und R^2 zusammen einen Rest der Formel $-N=CH-NH-$ oder $-NH-CO-NH-$ bilden, die Cyclisierung einer Verbindung der Formel I, wobei R^1 eine Aminogruppe ist und R^2 eine Aminogruppe, ein (1-4C)Alkylamino-, (3-4C)Alkenylamino-, (3-4C)Alkynylamino- oder ein substituiertes (1-4C)Alkylaminorest ist, mit einer geeigneten Carbonsäure, einem Amid einer Carbonsäure, einem Harnstoff oder einem Carbonat;
- (c) zur Herstellung jener gegebenenfalls substituierten Verbindungen der Formel I, wobei R^1 und R^2 zusammen einen Rest der Formel $-N=N-NH-$ bilden, die Diazotierung und Cyclisierung einer Verbindung der Formel I wobei R^1 eine Aminogruppe ist und R^2 eine Aminogruppe, ein (1-4C)Alkylamino-, (3-4C)Alkenylamino-, (3-4C)

Alkylamino- oder ein substituierter (1-4C)Alkylaminorest ist;

(d) zur Herstellung jener gegebenenfalls substituierten Verbindungen der Formel I, wobei R¹ und R² zusammen einen Rest der Formel -N=CH-CH=N- bilden, die Cyclisierung einer Verbindung der Formel I, wobei R¹ eine Aminogruppe ist und R² eine Aminogruppe ist, mit einem geeigneten Diketon;

(e) zur Herstellung jener gegebenenfalls substituierten Verbindungen der Formel I, wobei R¹ und R² zusammen einen Rest der Formel -N=CH-CO-NH- bilden, die Cyclisierung einer Verbindung der Formel I, wobei R¹ eine Aminogruppe ist und R² eine Aminogruppe, ein (1-4C)Alkylamino-, (3-4C)Alkenylamino-, (3-4C)Alkylamino- oder ein substituierter (1-4C)Alkylaminorest ist, mit einer geeigneten Dicarbonsäure oder einem Diester davon; oder

(f) zur Herstellung jener Verbindungen der Formel I, die einen (1-4C)Alkylsulfinyl- oder (1-4C)Alkylsulfonyl-substituenten tragen, die Oxidation eines Chinazolinderivats der Formel I, das einen (1-4C)Alkylthiosubstituenten trägt;


und wenn ein pharmazeutisch verträgliches Salz eines tricyclischen Derivats der Formel I erforderlich ist, kann es unter Verwendung eines üblichen Verfahrens erhalten werden.

10. Arzneimittel, das ein tricyclisches Derivat der Formel I oder ein pharmazeutisch verträgliches Salz davon umfaßt, wie in einem der Ansprüche 1 bis 8 beansprucht, zusammen mit einem pharmazeutisch verträglichen Verdünnungsmittel oder Träger.

11. Verwendung eines tricyclischen Derivats der Formel I oder eines pharmazeutisch verträglichen Salzes davon, wie in einem der Ansprüche 1 bis 8 beansprucht, bei der Herstellung eines Medikaments zur Verwendung bei der Erzeugung einer Antikrebswirkung in einem Warmblüter.

Revendications

1. Dérivé tricyclique de formule I




où R¹ et R² forment ensemble un groupe de formule -N=CH-NH-, -N=CH-O-, -N=CH-S-, -N=N-NH-, -NH-N=CH-, -NH-CH=CH-, -NH-CO-NH-, -NH-CO-O-, -NH-CO-S-, -NH-NH-CO-, -N=CH-CH=CH-, -N=N-CH=CH-, -N=CH-N=CH-, -N=CH-CH=N-, -NH-CO-CHCH- ou -N=CH-CO-NH- (un atome d'azote étant situé dans chaque cas en position 6 du cycle quinazoline) et le cycle à 5 ou 6 chaînons ainsi formé peut éventuellement porter un ou deux substituants, un substituant quelconque sur un atome d'azote disponible étant choisi parmi alkyle en C₁₋₄, alcényle en C₃₋₄, alcynyle en C₃₋₄, halogéno-alkyle en C₁₋₄, hydroxyalkyle en C₁₋₄, alcanoyloxy en C₂₋₄, alkyle en C₁₋₄, alcoxy en C₁₋₄, alkyle en C₁₋₄, cyanoalkyle en C₁₋₄, aminoalkyle en C₁₋₄, alkylamino en C₁₋₄, alkyle en C₁₋₄ et di-[alkyle en C₁₋₄]aminoalkyle en C₁₋₄, et un substituant quelconque sur un atome de carbone disponible étant choisi parmi halogéno, amino, hydroxyle, carbamoyle, cyano, alkyle en C₁₋₄, alcényle en C₂₋₄, alcynyle en C₂₋₄, alcoxy en C₁₋₄, alkylthio en C₁₋₄, alkylsulfinyle en C₁₋₄, alkylsulfonyle en C₁₋₄, alkylamino en C₁₋₄, di-[alkyle en C₁₋₄]amino, alcanoyle en C₂₋₄, N-alkyle en C₁₋₄-carbamoyle, N,N-di-[alkyle en C₁₋₄]carbamoyle, halogénoalkyle en C₁₋₄, hydroxyalkyle en C₁₋₄, alcanoyloxy en C₂₋₄, alkyle en C₁₋₄, alcoxy en C₁₋₄, alkyle en C₁₋₄, cyanoalkyle en C₁₋₄, aminoalkyle en C₁₋₄, alkylamino en C₁₋₄ et di-[alkyle en C₁₋₄]aminoalkyle en C₁₋₄; et m est l'entier 1, 2 ou 3 et chaque R³ est indépendamment hydrogène, halogéno, trifluorométhyle, hydroxyle, amino, nitro, cyano, alkyle en C₁₋₄, alcoxy en C₁₋₄, alkylamino en C₁₋₄, di-[alkyle en C₁₋₄]amino ou alcanoylamino en C₂₋₄; ou sel pharmaceutiquement acceptable de celui-ci.


2. Dérivé tricyclique de formule I selon la revendication 1 où le cycle tricyclique 6,6,5 défini par la liaison des groupes R¹ et R² est choisi parmi 3H-imidazo[4,5-g]quinazolin-8-yle, oxazo[4,5-g]quinazolin-8-yle, thiazolo[4,5-g]quinazolin-8-yle, 3H-[1,2,3]triazolo[4,5-g]quinazolin-8-yle, 1H-pyrazolo[3,4-g]quinazolin-8-yle, 6H-pyrrolo[2,3-g]quinazolin-4-yle, 2-oxo-1,2-dihydro-3H-imidazo[4,5-g]quinazolin-8-yle, 2-oxo-1,2-dihydrooxazo[4,5-g]quinazolin-8-yle, 2-oxo-1,2-dihydrothiazolo[4,5-g]quinazolin-8-yle et 3-oxo-2,3-dihydro-1H-pyrazolo[3,4-g]quinazolin-8-yle, et le cycle à 5 chaînons comprenant R¹ et R² peut éventuellement porter un ou deux substituants, un substituant quelconque sur un atome d'azote disponible étant choisi parmi méthyle, éthyle, propyle, allyle, prop-2-ynyle, 2,2,2-trifluoroéthyle, 2-hydroxyéthyle, 3-hydroxypropyle, 2-acétoxyéthyle, 2-méthoxyéthyle, 3-méthoxypropyle, cyanométhyle, 2-cyanoéthyle, 2-aminoéthyle, 2-méthylaminoéthyle et 2-diméthylaminoéthyle et un substituant quelconque sur un atome de carbone disponible étant choisi parmi fluoro, chloro, amino, carbamoyle, cyano, méthyle, éthyle, propyle, vinyle, allyle, éthynyle, prop-2-ynyle, méthoxy, éthoxy, propoxy, méthylthio, méthylsulfonyle, méthylsulfonyle, méthylamino, diméthylamino, acétyl, N-méthylcarbamoyle, N,N-diméthylcarbamoyle, fluorométhyle, difluorométhyle, trifluorométhyle, 2,2,2-trifluoroéthyle, hydroxyméthyle, 2-hydroxyéthyle, acétoxyméthyle, 2-acétoxyéthyle, méthoxyméthyle, 2-méthoxyéthyle, cyanométhyle, 2-cyanoéthyle, aminométhyle, 2-aminoéthyle, méthylaminométhyle, 2-méthylaminoéthyle, diméthylaminométhyle et 2-diméthylaminoéthyle ; et m est l'entier 1, 2 ou 3 et chaque R³ est indépendamment hydrogène, fluoro, chloro, bromo, trifluorométhyle, hydroxyle, amino, nitro, cyano, méthyle, éthyle, méthoxy, méthylamino, diméthylamino ou acétamido ; ou sel pharmaceutiquement acceptable de celui-ci.
3. Dérivé tricyclique de formule I selon la revendication 1, où le cycle tricyclique 6,6,6 défini par la liaison des groupes R¹ et R² est choisi parmi pyrido[2,3-g]quinazolin-4-yle, pyrimidino[4,5-g]cinnolin-9-yle, pyrimidino[4,5-g]quinazolin-4-yle, pyrazino[2,3-g]quinazolin-4-yle, 7-oxo-6,7-dihydropyrido[2,3-g]quinazolin-4-yle et 8-oxo-8,9-dihydropyrazino[2,3-g]quinazolin-4-yle, et le cycle à 6 chaînons comprenant R¹ et R² peut éventuellement porter un ou deux substituants, un substituant quelconque sur un atome d'azote disponible étant choisi parmi méthyle, éthyle et propyle, et un substituant quelconque sur un atome de carbone disponible étant choisi parmi fluoro, chloro, hydroxyle, carbamoyle, cyano, méthyle, méthoxy, éthoxy, N-méthylcarbamoyle, N,N-diméthylcarbamoyle, trifluorométhyle et 2,2,2-trifluoroéthyle ; et m est l'entier 1 ou 2 et chaque R³ est indépendamment hydrogène, fluoro, chloro, bromo, trifluorométhyle, nitro, cyano, méthyle ou éthyle ; ou sel pharmaceutiquement acceptable de celui-ci.
4. Dérivé tricyclique de formule I selon la revendication 1 où le cycle tricyclique 6,6,5 défini par la liaison des groupes R¹ et R² est choisi parmi 3-méthyl-3H-imidazo[4,5-g]quinazolin-8-yle, 3-méthyl-3H-[1,2,3]triazolo[4,5-g]quinazolin-8-yle et 3-méthyl-2-oxo-1,2-dihydro-3H-imidazo[4,5-g]quinazolin-8-yle ; et (R³)_m est 3'-méthyle, 3'-chloro ou 3'-chloro-4'-fluoro ; ou sel pharmaceutiquement acceptable de celui-ci.
5. Dérivé tricyclique de formule I selon la revendication 1, où le cycle tricyclique 6,6,5 défini par la liaison des groupes R¹ et R² est choisi parmi 3-méthyl-3H-imidazo[4,5-g]quinazolin-8-yle, 3-méthyl-2-oxo-1,2-dihydro-3H-imidazo[4,5-g]quinazolin-8-yle et 3-méthyl-2-trifluorométhyl-3H-imidazo[4,5-g]quinazolin-8-yle ; et (R³)_m est 3'-méthyle, 3'-chloro ou 3'-chloro-4'-fluoro ; ou sel pharmaceutiquement acceptable de celui-ci.
6. Dérivé tricyclique de formule I selon la revendication 1 où le cycle tricyclique 6,6,6 défini par la liaison des groupes R¹ et R² est choisi parmi 7,8-diméthylpyrazino[2,3-g]quinazolin-4-yle et 7-hydroxy-9-méthyl-8-oxo-8,9-dihydropyrazino[2,3-g]quinazolin-4-yle ; et (R³)_m est 3'-méthyle, 3'-chloro ou 3'-chloro-4'-fluoro ; ou sel pharmaceutiquement acceptable de celui-ci.
7. Dérivé tricyclique de formule I selon la revendication 1, choisi parmi : la 3-méthyl-8-(3'-méthylanilino)-3H-imidazo[4,5-g]quinazoline, la 3-méthyl-8-(3'-méthylanilino)-1,2-dihydro-3H-imidazo[4,5-g]quinazolin-2-one ou la 8-(3'-chloro-4'-fluoroanilino)-3-méthyl-2-trifluorométhyl-3H-imidazo[4,5-g]quinazoline ; ou sel pharmaceutiquement acceptable de celui-ci.
8. Dérivé tricyclique de formule I selon la revendication 1, choisi parmi : la 4-(3'-chloro-4'-fluoroanilino)-7,8-diméthylpyrazino[2,3-g]quinazoline ou la 4-(3'-chloro-4'-fluoroanilino)-7-hydroxy-9-méthyl-8-oxo-8,9-dihydropyrazino[2,3-g]quinazoline ; ou sel pharmaceutiquement acceptable de celui-ci.

9. Procédé pour la préparation d'un dérivé tricyclique de formule I, ou d'un sel pharmaceutiquement acceptable de celui-ci, selon la revendication 1, qui comprend :

(a) la réaction d'une quinazoline de formule II



où Z est un groupe déplaçable, avec une aniline de formule III



- (b) pour la production des composés éventuellement substitués de formule I où R¹ et R² forment ensemble un groupe de formule -N=CH-NH- ou -NH-CO-NH-, la cyclisation d'un composé de formule I où R¹ est amino et R² est amino, alkylamino en C₁₋₄, alcénylamino en C₃₋₄, alcynylamino en C₃₋₄ ou un alkylamino en C₁₋₄ substitué avec un acide carboxylique approprié, un amide d'un acide carboxylique, une urée ou un carbonate ;
- (c) pour la production des composés éventuellement substitués de formule I où R¹ et R² forment ensemble un groupe de formule -N=N-NH-, la diazotation et la cyclisation d'un composé de formule I où R¹ est amino et R² est amino, alkylamino en C₁₋₄, alcénylamino en C₃₋₄, alcynylamino en C₃₋₄ ou un alkylamino en C₁₋₄ substitué ;
- (d) pour la production des composés éventuellement substitués de formule I où R¹ et R² forment ensemble un groupe de formule -N=CH-CH=N-, la cyclisation d'un composé de formule I où R¹ est amino et R² est amino avec une dicétone appropriée ;
- (e) pour la production des composés éventuellement substitués de formule I où R¹ et R² forment ensemble un groupe de formule -N=CH-CO-NH-, la cyclisation d'un composé de formule I où R¹ est amino et R² est amino, alkylamino en C₁₋₄, alcénylamino en C₃₋₄, alcynylamino en C₃₋₄ ou un alkylamino en C₁₋₄ substitué avec un acide dicarboxylique approprié ou un diester de celui-ci ; ou
- (f) pour la production des composés de formule I qui portent un substituant alkylsulfinyle en C₁₋₄ ou alkylsulfonyl en C₁₋₄, l'oxydation d'un dérivé de quinazoline de formule I qui porte un substituant alkylthio en C₁₋₄ ;

et quand un sel pharmaceutiquement acceptable d'un dérivé tricyclique de formule I est nécessaire, il peut être obtenu au moyen d'un protocole conventionnel.

10. Composition pharmaceutique qui comprend un dérivé tricyclique de formule I, ou un sel pharmaceutiquement acceptable de celui-ci, selon l'une quelconque des revendications 1 à 8 en association avec un diluant ou support pharmaceutiquement acceptable.
11. Utilisation d'un dérivé tricyclique de formule I, ou d'un sel pharmaceutiquement acceptable de celui-ci, selon l'une quelconque des revendications 1 à 8 dans la fabrication d'un médicament destiné à être utilisé dans la production d'un effet anticancéreux chez un animal à sang chaud.